

polyéthylène, les microsphères de copolymères telles que l'Expancel (NOBEL INDUSTRIE), le Polytrap (DOW CORNING) et les microbilles de résine de silicone (Tospearl de TOSHIBA, par exemple).

- 5 Les nacres peuvent être présentes dans l'émulsion à raison de 0 à 20 % en poids, par rapport au poids total de l'émulsion, de préférence de 2 à 15 %.

Les nacres utilisables selon l'invention sont par exemple le mica recouvert d'oxyde de titane, d'oxyde de fer, de pigment naturel ou d'oxychlorure de bismuth, ainsi que le mica titane coloré.

10

L'émulsion selon l'invention peut comprendre en outre tous additifs usuellement utilisés dans le domaine cosmétique, tels que des antioxydants, des colorants, des parfums, des huiles essentielles, des conservateurs, des actifs cosmétiques, des hydratants, des vitamines, des sphingolipides, des filtres solaires, des polymères liposolubles, notamment hydrocarbonés, tels que le polybutène, les polyalkylènes, les polyacrylates et les polymères siliconés compatibles avec les corps gras. Bien entendu, l'homme du métier

15

20

quantité, de manière telle que les propriétés avantageuses de la composition selon l'invention ne soient pas, ou substantiellement pas, altérées par l'adjonction envisagée. Ces additifs peuvent être présents dans la composition à raison de 0 à 15% en poids.

- 25 Les émulsions selon l'invention peuvent se présenter sous la forme d'un produit cosmétique et notamment sous la forme d'un produit de maquillage, en particulier un fond de teint, un fard à joues ou à paupières, ou un rouge à lèvres.

- 30 Elles peuvent également se présenter sous forme non colorée, contenant éventuellement des actifs cosmétiques.

Les exemples qui suivent servent à illustrer l'invention sans toutefois présenter un caractère limitatif. Dans ces exemples, les quantités sont données en pourcentage de poids par rapport au poids total de la composition.

5

EXEMPLE 1 :

La Demanderesse a réalisé le fond de teint suivant :

10

PHASE H

	- Huile de jojoba hydrogénée	5,6 %
	- Conservateur	0,3 %
15	- Cire de polyéthylène (PM : 500)	2,9 %
	- Cire de polytétrafluoroéthylène	7 %
	- Mélange de poly méthylcétyl diméthyl méthyl- siloxane oxyéthyléné oxypropyléné, isostéarate polyglycérolé (4 moles) et laurate d'hexyle,	
20	vendu sous la dénomination "ABIL WE 09" par la Société GOLDSCHMIDT	9 %

SILICONE S

25	- Cyclohexadiméthylsiloxane (viscosité : 8.10^{-6} m ² /s)	24,3 %
----	-------------------------------------------------------------------------	--------

PIGMENTS P

	- Oxydes de fer	2,9 %
30	- Oxyde de titane	7,1 %

PHASE E

	- Eau déminéralisée stérilisée	38,5 %
	- Propylèneglycol	1 %
	- Conservateur	0,4 %
5	- Sulfate de magnésium, 7 H ₂ O	1 %

On pèse ensemble les composés de la phase H que l'on chauffe à 100°C.

Après homogénéisation, on la refroidit à 80 °C puis on additionne la silicone S.

10 Les pigments P sont ensuite dispersés dans le mélange H+S.

Après homogénéisation, on ajoute lentement la phase E, préalablement chauffée à 80 °C, en agitant à l'aide d'un agitateur type Moritz et en conservant au cours de l'addition une température minimale de 75 °C.

On coule le produit et on obtient un fond de teint compact homogène, qui
15 s'étale bien, qui conduit à un maquillage très naturel et très doux, qui présente une bonne tenue cosmétique et qui ne transfère pas.

Cette composition a une force de pénétration, mesurée sur un appareil STEVENS comme décrit dans le texte ci-dessus, supérieure à 50 grammes.

20

EXEMPLE 2 : COMPARATIF : Fond de teint selon l'Art antérieur

On a reproduit l'exemple 5 de la demande JP-A-03261707.

25

PHASE H

	- Paraffine (cire minérale)	4 %
	- Mélange diméthicone copolyol, cyclopentasiloxane,	
30	eau (10/88/2), vendu sous la dénomination "Q2-3225 C"	
	par la Société DOW CORNING	30 %
	- Camphre	0,1 %

- Menthol	0,1	%
- Cellulose microcristalline	3	%

PIGMENTS P

5

- Dioxyde de titane enrobé de polydiméthylsiloxane	15	%
- Oxyde de fer jaune enrobé de polydiméthylsiloxane	3	%
- Oxyde de fer rouge enrobé de polydiméthylsiloxane	1	%
- Oxyde de fer noir enrobé de polydiméthylsiloxane	0,2	%

10

PHASE E

- Eau	33	%
15 - Ethanol	5	%
- Polyéthylèneglycol	5	%
- p-hydroxybenzoate de méthyle	0,3	%

On prépare un fond de teint compact selon le mode opératoire ci-dessus.

- 20 On obtient un fond de teint de couleur hétérogène, qui est inhomogène, dont les pigments sont mal dispersés et qui est rêche et sec à l'application.

25 EXEMPLE 3 :

La Demanderesse a réalisé les trois compositions A, B, et C suivantes, en faisant varier la nature de la cire :

30 PHASE H

- cire	6,3	%
--------	-----	---

WO 99/47111

PCT/FR99/00609

	- conservateur	0,3 %
	- Mélange de poly méthylcétyl diméthyl méthyl- siloxane oxyéthyléné oxypropyléné, isostéarate polyglycérolé (4 moles) et laurate d'hexyle,	
5	vendu sous la dénomination "ABIL WE 09" par la Société GOLDSCHMIDT	9 %

SILICONE S

10	- Cyclohexadiméthylsiloxane (viscosité : $8 \cdot 10^{-6}$ m ² /s)	24,3 %
----	-------------------------------------------------------------------------------	--------

PIGMENTS P

	- Oxydes de fer	2,9 %
15	- Oxyde de titane	7,1 %

PHASE E

	- Eau déminéralisée stérilisée	38,5 %
20	- Propylèneglycol	1 %
	- Conservateur	0,4 %
	- Sulfate de magnésium, 7 H ₂ O	1 %

avec :

25

30

Composition	A	B	C
Nature de la cire	huile de jojoba hydrogénée	polyéthylène	carnauba
Force de pénétration (en g)	74	101	11
Aspect après coulage à chaud	solide non lisse	solide	mou
Observation au microscope	émulsion fine mais pas très régulière	émulsion fine régulière	petits cristaux


Ces trois émulsions ont été préparées comme dans l'exemple 1. La force de pénétration a été mesurée sur un appareil STEVENS comme décrit dans le texte ci-dessus.

5

L'émulsion C qui ne comprend que de la cire de carnauba ne permet pas d'obtenir une composition suffisamment solide.

REVENDICATIONS

1. Utilisation d'un tensio-actif siliconé de type alkyl-diméthicone copolyol de
5 formule :



dans laquelle :

- 20 PE = $(-\text{C}_2\text{H}_4\text{O})_x (-\text{C}_3\text{H}_6\text{O})_y - \text{H}$,
 $x = 0$ à 50 ,
 $y = 0$ à 30 , x et y n'étant pas simultanément 0 ,
 $o = 1$ à 100 ,
 $m = 1$ à 40 ,
25 $n = 1$ à 200 ,
 $p = 1$ à 17 et
 $q = 1$ à 5 .

- pour la préparation d'émulsions solides eau-dans-huile, comportant une phase
30 aqueuse émulsionnée par ledit tensio-actif dans une phase grasse comportant
au moins une huile et au moins une cire.

2. Utilisation selon la revendication 1, caractérisée par le fait que :

- o = 1 à 25,
5 m = 1 à 10,
n = 1 à 100.

3. Utilisation selon la revendication 2, caractérisée par le fait que :

- 10 o = 21,
m = 4,
n = 73,

4. Utilisation selon la revendication 3, caractérisée par le fait que le tensio-actif
15 siliconé est un mélange de cétyldiméthicone copolyol, de polyglycérol-4 isostéarate et de laurate d'hexyle.

5. Utilisation selon l'une quelconque des revendications 1 à 4, caractérisée par
le fait que le tensio-actif siliconé alkyldiméthicone copolyol est utilisé à raison
20 de 0,5 à 40% et de préférence 2 à 12% en poids, par rapport au poids total de l'émulsion.

6. Utilisation selon l'une quelconque des revendications 1 à 5, caractérisée par
le fait que la phase grasse comprend une huile de silicone.

25

7. Utilisation selon la revendication 6, caractérisée par le fait que l'huile de
silicone est choisie parmi les silicones volatiles cycliques ayant de 3 à 8
atomes de silicium, les silicones volatiles linéaires ayant de 2 à 9 atomes de
silicium, les cyclocopolymères diméthylsiloxane/ méthylalkyl-siloxane, les
30 polyalkylsiloxanes à groupements terminaux triméthyl-silyle et les huiles de
silicone phénylées.

8. Utilisation selon la revendication 6 ou 7, caractérisée par le fait que l'huile de silicone est une huile de silicone volatile.
9. Utilisation selon la revendication 8, caractérisée par le fait que l'huile de silicone volatile est choisie parmi le cyclotétradiméthyl-siloxane, le cyclopentadiméthylsiloxane, le cyclohexadiméthyl-siloxane, l'hexaméthylidisiloxane, l'hexylheptaméthyltrisiloxane et l'octylheptaméthyltrisiloxane.
10. Utilisation selon l'une quelconque des revendications 1 à 7, caractérisée par le fait que la phase grasse comprend une isoparaffine volatile.
11. Utilisation selon la revendication 10, caractérisée par le fait que l'isoparaffine volatile est une isoparaffine en C8-C16 choisie parmi l'isododécane, l'isodécane et l'isohexadécane.
12. Utilisation selon l'une quelconque des revendications 1 à 11, caractérisée par le fait que la phase grasse de l'émulsion solide eau-dans-huile comprend en outre des huiles minérales, des huiles d'origine animale, des huiles végétales, des esters ramifiés en C8-C16, des esters et éthers de synthèse, des esters hydroxylés, des esters de polyols, des alcools gras, des huiles fluorées et leurs mélanges.
13. Utilisation selon l'une quelconque des revendications 1 à 12, caractérisée par le fait que les huiles sont utilisées à raison de 10 à 40% et de préférence de 18 à 30 % en poids, par rapport au poids total de l'émulsion.
14. Utilisation selon l'une quelconque des revendications 1 à 13, caractérisée par le fait que la phase grasse comprend au moins une cire ou un mélange de cires apte à conférer à l'émulsion une force de pénétration supérieure ou égale à 50 grammes.

15. Utilisation selon l'une quelconque des revendications 1 à 14 caractérisée par le fait que la cire est choisie parmi les cires végétales, minérales, animales, synthétiques, les huiles hydrogénées concrètes à 25°C, les esters gras concrets à 25°C et leurs mélanges.
- 5
16. Utilisation selon l'une quelconque des revendications 1 à 15, caractérisée par le fait que la cire est choisie parmi la cire de polyéthylène, l'huile de jojoba hydrogénée, l'ozokérite ou leurs mélanges.
- 10
17. Utilisation selon l'une quelconque des revendications 1 à 16, caractérisée par le fait que la cire est un mélange de cire de polyéthylène et d'huile de jojoba hydrogénée.
18. Utilisation selon l'une quelconque des revendications 1 à 14, caractérisée
- 15 par le fait que les cires sont utilisées à raison de 3 à 15 % et de préférence de 3 à 10 % en poids, par rapport au poids total de l'émulsion.
19. Utilisation selon l'une quelconque des revendications 1 à 18, caractérisée par le fait que la phase grasse de l'émulsion solide eau-dans-huile contient en
- 20 outre des pigments et/ou des nacres et/ou des charges sélectionnés pour un transfert minimum.
20. Utilisation selon l'une quelconque des revendications 1 à 19, caractérisée par le fait que la phase aqueuse représente 0,5 à 60 % du poids total de
- 25 l'émulsion.
21. Utilisation selon l'une quelconque des revendications 1 à 20, caractérisée par le fait que la phase aqueuse comprend de l'eau ou une eau florale, 0 à 14 % en poids par rapport au poids total de la phase aqueuse de monoalcools
- 30 inférieurs en C2-C6 et/ou de polyols et 0 à 6% en poids par rapport au poids total de l'émulsion d'un agent épaississant.


22. Utilisation selon l'une quelconque des revendications 1 à 21, d'un tensio-actif siliconé du type alkylidiméthicone copolyol de formule indiquée à l'une des revendications 1 à 4, pour l'obtention d'un produit de maquillage et notamment d'un fond de teint compact sans transfert.

5

23. Emulsion cosmétique solide, du type eau-dans-huile, caractérisée par le fait qu'elle comporte une phase aqueuse émulsionnée à l'aide d'un tensio-actif siliconé de type alkylidiméthicone copolyol de formule :

10

15



20

dans laquelle :

- 25 PE = $(-\text{C}_2\text{H}_4\text{O})_x (-\text{C}_3\text{H}_6\text{O})_y - \text{H}$,
 $x = 0 \text{ à } 50$,
 $y = 0 \text{ à } 30$, x et y n'étant pas simultanément 0,
 $o = 1 \text{ à } 100$,
 $m = 1 \text{ à } 40$,
30 $n = 1 \text{ à } 200$,
 $p = 1 \text{ à } 17$ et
 $q = 1 \text{ à } 5$.

dans une phase grasse contenant au moins une huile et au moins une cire.

24. Emulsion selon la revendication 23, caractérisée par le fait que :

5

$o = 1$ à 25,
 $m = 1$ à 10,
 $n = 1$ à 100.

10 25. Emulsion selon la revendication 24, caractérisée par le fait que :

$o = 21$,
 $m = 4$,
 $n = 73$,

15

26. Emulsion selon la revendication 25, caractérisée par le fait que le tensio-actif siliconé est un mélange de cétyldiméthicone copolyol, de polyglycéryl-4 isostéarate et de laurate d'hexyle.

20 27. Emulsion selon l'une quelconque des revendications 23 à 26, caractérisée par le fait qu'elle contient 0,5 à 40 % et de préférence 2 à 12 % en poids de tensio-actif siliconé alkyldiméthicone copolyol, sur la base du poids total de l'émulsion.

25 28. Emulsion selon l'une quelconque des revendications 23 à 27, caractérisée par le fait que la phase grasse comprend une huile de silicone.

29. Emulsion selon la revendication 28, caractérisée par le fait que l'huile de silicone est choisie parmi les silicones volatiles cycliques ayant de 3 à 8
30 atomes de silicium, les silicones volatiles linéaires ayant de 2 à 9 atomes de silicium, les cyclocopolymères diméthylsiloxane/ méthylalkylsiloxane, les polyalkylsiloxanes à groupements terminaux triméthyl-silyle et les huiles de

silicone phénylées.

30. Emulsion selon la revendication 28 ou 29, caractérisée par le fait que l'huile de silicone est une huile de silicone volatile.

5

31. Emulsion selon la revendication 30, caractérisée par le fait que l'huile de silicone volatile est choisie parmi le cyclotétradiméthyl-siloxane, le cyclopentadiméthylsiloxane, le cyclohexadiméthylsiloxane, l'hexaméthylidisiloxane, l'hexylhepta-méthyltrisiloxane et l'octylhepta-méthyltrisiloxane.

10

32. Emulsion selon l'une quelconque des revendications 23 à 27, caractérisée par le fait que la phase grasse comprend une isoparaffine volatile.

15 33. Emulsion selon la revendication 32, caractérisée par le fait que l'isoparaffine volatile est une isoparaffine en C8-C16 choisie parmi l'isododécane, l'isodécane et l'isohexadécane.

20 34. Emulsion selon l'une quelconque des revendications 23 à 33, caractérisée par le fait que la phase grasse comprend en outre des huiles minérales, des huiles d'origine animale, des huiles végétales, des esters ramifiés en C8-C16, des esters et éthers de synthèse, des esters hydroxylés, des esters de polyols, des alcools gras, des huiles fluorées et leurs mélanges.

25 35. Emulsion selon l'une quelconque des revendications 23 à 34, caractérisée par le fait qu'elle contient 10 à 40 % et de préférence 18 à 30 % en poids d'huile(s), sur la base du poids total de l'émulsion.

30 36. Emulsion selon l'une quelconque des revendications 23 à 35, caractérisée par le fait que la phase grasse comprend au moins une cire ou un mélange de cires apte à conférer à l'émulsion une force de pénétration supérieure ou égale à 50 grammes.

37. Emulsion selon l'une quelconque des revendications 23 à 36, caractérisée par le fait que la cire est choisie parmi les cires minérales, animales, végétales, synthétiques, les huiles hydrogénées concrètes à 25°C, les esters gras concrets à 25°C et leurs mélanges.
38. Emulsion selon l'une quelconque des revendications 23 à 37, caractérisée par le fait que la cire est choisie parmi la cire de polyéthylène, l'huile de jojoba hydrogénée, l'ozokérite ou leurs mélanges.
39. Emulsion selon l'une quelconque des revendications 23 à 38, caractérisée par le fait que la cire est un mélange de cire de polyéthylène et d'huile de jojoba hydrogénée.
40. Emulsion selon l'une quelconque des revendications 23 à 39, caractérisée par le fait qu'elle contient 3 à 15 % et de préférence 3 à 10 % en poids de cire(s), sur la base du poids total de l'émulsion.
41. Emulsion selon l'une quelconque des revendications 23 à 40, caractérisée par le fait que la phase grasse comprend en outre des pigments et/ou des nacres et/ou des charges sélectionnés pour un transfert minimum.
42. Emulsion selon l'une quelconque des revendications 23 à 41 caractérisée par le fait que la phase aqueuse représente 0,5 à 60 % du poids total de l'émulsion.
43. Emulsion selon l'une quelconque des revendications 23 à 42, caractérisée par le fait que la phase aqueuse comprend de l'eau ou une eau florale, 0 à 14 % en poids par rapport au poids total de la phase aqueuse, de monoalcools inférieurs en C2-C6 et/ou de polyols et 0 à 6 % en poids par rapport au poids total de l'émulsion d'un agent épaississant, ainsi qu'éventuellement des agents de stabilisation de l'émulsion.

44. Emulsion selon l'une quelconque des revendications 23 à 43, caractérisée par le fait qu'elle comprend en outre au moins un additif choisi parmi les antioxydants, les colorants, les parfums, les huiles essentielles, les conservateurs, les actifs cosmétiques, les hydratants, les vitamines, les sphingolipides, les filtres solaires et les polymères liposolubles.
45. Emulsion selon l'une quelconque des revendications 23 à 44, caractérisée par le fait qu'elle est constituée par un produit de maquillage, et notamment un fond de teint compact sans transfert.
46. Procédé de maquillage de la peau et/ou du cuir chevelu, caractérisé par le fait qu'il consiste à appliquer sur la peau et/ou le cuir chevelu une émulsion solide telle que définie dans l'une quelconque des revendications 23 à 45.

INTERNATIONAL SEARCH REPORT

In ternational Application No

PCT/FR 99/00609

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K7/48 A61K7/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>EP 0 595 683 A (OREAL) 4 May 1994 (1994-05-04) page 6, line 26-58 page 7, line 4-9 page 7, line 40-58 page 8, column -12 page 8, line 48-58 page 9, line 31,32 - line 47-49 examples 29,32</p> <p style="text-align: center;">--- -/--</p>	<p>1-9, 12-46</p>

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Z" document member of the same patent family

Date of the actual completion of the international search

16 July 1999

Date of mailing of the international search report

27/07/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.
Fax: (+31-70) 340-3016

Authorized officer

Sierra Gonzalez, M

INTERNATIONAL SEARCH REPORT

International Application No

PCT/FR 99/00609

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>D. G. KRZYSIK: "A New Silicone Emulsifier For Water-in-Oil Systems"</p> <p>DRUG COSMET. IND.,</p> <p>vol. 146, no. 4, 1990, XP002090584</p> <p>pages 28, 30, 35, 79, 81</p> <p>page 30, left-hand column, paragraph 4</p> <p>page 35, left-hand column, last last</p> <p>page 79, left-hand column, last paragraph</p> <p>---</p>	<p>1-3,5,</p> <p>12-15,</p> <p>19-23,</p> <p>34-37,</p> <p>41-46</p>
A	<p>F. BATTIONI: "Dalle emulsioni A/O alle emulsioni A/S e viceversa"</p> <p>IL PRODOTTO CHIMICO,</p> <p>vol. 28, no. 3, 1987, page 32-34</p> <p>XP002090585</p> <p>the whole document</p> <p>---</p>	<p>1-46</p>
A	<p>EP 0 374 332 A (SHISEIDO CO LTD)</p> <p>27 June 1990 (1990-06-27)</p> <p>the whole document</p> <p>---</p>	<p>1-46</p>
A	<p>CHEMICAL ABSTRACTS, vol. 116, no. 10,</p> <p>9 March 1992 (1992-03-09)</p> <p>Columbus, Ohio, US;</p> <p>abstract no. 91163,</p> <p>TAKADA, SUSUMU ET AL: "Solid-type water-in-oil emulsion cosmetics containing silicone oils, waxes, and spherical powders"</p> <p>XP002090586</p> <p>cited in the application</p> <p>abstract</p> <p>& JP 03 261707 A (SHISEIDO CO., LTD., JAPAN)</p> <p>-----</p>	<p>1-46</p>

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/FR 99/00609

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0595683 A	04-05-1994	FR 2696931 A AT 152604 T CA 2108912 A DE 69310477 D DE 69310477 T ES 2101266 T JP 6233928 A US 5851539 A	22-04-1994 15-05-1997 22-04-1994 12-06-1997 21-08-1997 01-07-1997 23-08-1994 22-12-1998
EP 0374332 A	27-06-1990	DE 3877967 A JP 1079104 A JP 2691729 B US 5362482 A	11-03-1993 24-03-1989 17-12-1997 08-11-1993
JP 3261707 A	21-11-1991	NONE	

RAPPORT DE RECHERCHE INTERNATIONALE

De de internationale No
PCT/FR 99/00609

A. CLASSEMENT DE L'OBJET DE LA DEMANDE
CIB 6 A61K7/48 A61K7/06

Selon la classification internationale des brevets (CIB) ou à la fois selon la classification nationale et la CIB

B. DOMAINES SUR LESQUELS LA RECHERCHE A PORTE

Documentation minimale consultée (système de classification suivi des symboles de classement)
CIB 6 A61K

Documentation consultée autre que la documentation minimale dans la mesure où ces documents relèvent des domaines sur lesquels a porté la recherche

Base de données électronique consultée au cours de la recherche internationale (nom de la base de données, et si réalisable, termes de recherche utilisés)

C. DOCUMENTS CONSIDERES COMME PERTINENTS

Catégorie *	Identification des documents cités, avec, le cas échéant, l'indication des passages pertinents	no. des revendications visées
X	<p>EP 0 595 683 A (OREAL) 4 mai 1994 (1994-05-04) page 6, ligne 26-58 page 7, ligne 4-9 page 7, ligne 40-58 page 8, colonne -12 page 8, ligne 48-58 page 9, ligne 31,32 - ligne 47-49 exemples 29,32</p> <p style="text-align: center;">--- -/--</p>	1-9, 12-46

☒ Voir la suite du cadre C pour la fin de la liste des documents

☒ Les documents de familles de brevets sont indiqués en annexe

* Catégories spéciales de documents cités:

- "A" document définissant l'état général de la technique, non considéré comme particulièrement pertinent
- "E" document antérieur, mais publié à la date de dépôt international ou après cette date
- "L" document pouvant jeter un doute sur une revendication de priorité ou cité pour déterminer la date de publication d'une autre citation ou pour une raison spéciale (telle qu'indiquée)
- "O" document se référant à une divulgation orale, à un usage, à une exposition ou tous autres moyens
- "P" document publié avant la date de dépôt international, mais postérieurement à la date de priorité revendiquée

- "T" document ultérieur publié après la date de dépôt international ou la date de priorité et n'appartenant pas à l'état de la technique pertinent, mais cité pour comprendre le principe ou la théorie constituant la base de l'invention
- "X" document particulièrement pertinent; l'invention revendiquée ne peut être considérée comme nouvelle ou comme impliquant une activité inventive par rapport au document considéré isolément
- "Y" document particulièrement pertinent; l'invention revendiquée ne peut être considérée comme impliquant une activité inventive lorsque le document est associé à un ou plusieurs autres documents de même nature, cette combinaison étant évidente pour une personne du métier
- "&" document qui fait partie de la même famille de brevets

Date à laquelle la recherche internationale a été effectivement achevée

16 juillet 1999

Date d'expiration du présent rapport de recherche internationale

27/07/1999

Nom et adresse postale de l'administration chargée de la recherche internationale
Office Européen des Brevets, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Fonctionnaire autorisé

Sierra Gonzalez, M

RAPPORT DE RECHERCHE INTERNATIONALE

De l'Organisation Mondiale de la Propriété Industrielle

PCT/FR 99/00609

C.(suite) DOCUMENTS CONSIDERES COMME PERTINENTS

Catégorie	Identification des documents cités, avec, le cas échéant, l'indication des passages pertinents	no. des revendications visées
X	<p>D. G. KRZYSIK: "A New Silicone Emulsifier For Water-in-Oil Systems" DRUG COSMET. IND., vol. 146, no. 4, 1990, XP002090584</p> <p>pages 28, 30, 35, 79, 81 page 30, colonne de gauche, alinéa 4 page 35, colonne de gauche, dernière dernière page 79, colonne de gauche, dernier alinéa</p> <p>---</p>	<p>1-3,5, 12-15, 19-23, 34-37, 41-46</p>
A	<p>F. BATTIONI: "Dalle emulsioni A/O alle emulsioni A/S e viceversa" IL PRODOTTO CHIMICO, vol. 28, no. 3, 1987, page 32-34 XP002090585 le document en entier</p> <p>---</p>	<p>1-46</p>
A	<p>EP 0 374 332 A (SHISEIDO CO LTD) 27 juin 1990 (1990-06-27) le document en entier</p> <p>---</p>	<p>1-46</p>
A	<p>CHEMICAL ABSTRACTS, vol. 116, no. 10, 9 mars 1992 (1992-03-09) Columbus, Ohio, US; abstract no. 91163, TAKADA, SUSUMU ET AL: "Solid-type water-in-oil emulsion cosmetics containing silicone oils, waxes, and spherical powders" XP002090586 cité dans la demande abrégé & JP 03 261707 A (SHISEIDO CO., LTD., JAPAN)</p> <p>-----</p>	<p>1-46</p>

RAPPORT DE RECHERCHE INTERNATIONALE

Renseignements relatifs aux membres de familles de brevets

C .nde internationale No

PCT/FR 99/00609

Document brevet cité au rapport de recherche	Date de publication	Membre(s) de la famille de brevet(s)	Date de publication
EP 0595683 A	04-05-1994	FR 2696931 A	22-04-1994
		AT 152604 T	15-05-1997
		CA 2108912 A	22-04-1994
		DE 69310477 D	12-06-1997
		DE 69310477 T	21-08-1997
		ES 2101266 T	01-07-1997
		JP 6233928 A	23-08-1994
		US 5851539 A	22-12-1998
EP 0374332 A	27-06-1990	DE 3877967 A	11-03-1993
		JP 1079104 A	24-03-1989
		JP 2691729 B	17-12-1997
		US 5362482 A	08-11-1993
JP 3261707 A	21-11-1991	AUCUN	

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 7/48, 7/42, 7/021	A1	(11) International Publication Number: WO 99/30681 (43) International Publication Date: 24 June 1999 (24.06.99)
(21) International Application Number: PCT/US98/26580 (22) International Filing Date: 15 December 1998 (15.12.98) (30) Priority Data: 08/990,460 15 December 1997 (15.12.97) US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 08/990,460 (CON) Filed on 15 December 1997 (15.12.97) (71) Applicant (for all designated States except US): NEOSTRATA COMPANY, INC. [US/US]; 4 Research Way, Princeton, NJ 08540 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): TSENG, Chung-Ye [US/US]; 43 Christy Drive, Warren, NJ 07059 (US). NAIK-SATAM, Prakash, G. [US/US]; 4 Lexington Avenue, Bloomfield, NJ 07003 (US). YEDNAK, Marie, E. [US/US]; 16 Half Acre Road, Jamesburg, NJ 08831 (US).		(74) Agents: SANDERCOCK, Colin, G. et al.; Foley & Lardner, Suite 500, 3000 K Street, N.W., Washington, DC 20007-5109 (US). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: COSMETIC MAKEUP COMPOSITION (57) Abstract <p>A water-in-silicone cosmetic makeup composition includes a hydroxy acid, a sunscreen agent, a moisturizing agent, an antioxidant, a cosmetically acceptable pigment, an emulsifier, a silicone-containing compound and water. The composition when applied to the skin protects the skin from ultraviolet light exposure, and it reduces the appearance of fine lines and wrinkles. The composition also is highly moisturizing and gentle enough for sensitive skin. Methods of protecting the skin from ultraviolet light exposure and reducing the appearance of fine lines and wrinkles by topically applying this composition, as well as methods of making the emulsified compositions also are disclosed.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

COSMETIC MAKEUP COMPOSITION

BACKGROUND OF THE INVENTION

1. Field of the Invention

5 This application relates to a makeup composition including a water in silicone emulsion containing hydroxyacids, such as alpha hydroxyacids, beta hydroxyacids and/or polyhydroxyacids. The makeup composition, when applied topically to the skin, provides beneficial effects to the skin such as (i) reducing the appearance of lines and wrinkles, (ii) protecting the skin from the harmful effects of the ultraviolet light from the sun, (iii) moisturizing and soothing the skin, and
10 (iv) making the skin attractively colored. This application also relates to methods of topical treatment of the skin to effect such benefits.

2. Description of the Prior Art

15 Human skin is comprised of two principal components, the outer epidermis and the underlying dermis, which is situated above the subcutaneous adipose or fat tissues. The epidermis consists of four distinct layers: stratum corneum, stratum granulosum, stratum spinosum and stratum basale; in the skin of palms and soles only, there is normally one additional zone called the stratum lucidum between the stratum corneum and the stratum granulosum.

20 The stratum corneum or horny layer is comprised of numerous flattened dead cells called corneocytes. The granular layer, located below the stratum corneum, contains large granules of keratinous materials. The spinous layer or prickle cell layer is located below the granular layer and composed of keratinocytes. The basal cell layer, located below the spinous layer but above the
25 dermis, is the only principal layer in the epidermis in which living cells normally replicate DNA and divide into daughter cells.

After a basal cell divides, one of the daughter cells migrates into the spinous layer where the cell starts to differentiate as a keratinocyte, wherein synthesis of keratin is initiated. As the keratinocyte continues to move outward
30 and reaches the granular layer, more keratinous materials are synthesized as

keratin filaments and large granules. The keratinocyte loses its nucleus in transit through the granular layer and thereupon dies to become a corneocyte without nucleus in the stratum corneum.

5 Normally, the granular layer is a few cell layers in thickness, and the stratum corneum may be 14-25 cell layers thick. Under normal conditions and in most skin areas, a keratinocyte takes 14 days to move outward from the basal cell layer to the granular layer, and a corneocyte takes 14 days to reach the outermost layer of the stratum corneum to be shed; the total time from basal cell layer to the surface is approximately 28 days. The entire sequence of epidermal terminal
10 differentiation is called keratinization.

The dermis is comprised mainly of collagen, elastic fibers, glycosaminoglycans and proteoglycans including hyaluronic acid, dermatan sulfate and chondroitin sulfate formerly known as mucopolysaccharides. Fibroblasts, the predominant cells of the dermis, synthesize collagen, elastic fibers, proteoglycans
15 and glycosaminoglycans. Collagen makes up approximately 77%, elastic fibers account for about 2%, and glycosaminoglycans constitute around 0.2% of the dry weight of the dermis. Collagen provides the tensile strength of and elastic fibers give resilience to the dermis. The glycosaminoglycans binds water to form a gelatinous mass between collagen and elastic fibers, which acts as a lubricant and
20 shock absorber for the dermis during movement of the skin.

Several factors affect human skin and result cutaneous aging to form wrinkles, sagging and loss of firmness and elasticity. These factors include internal factors and external factors. Cutaneous aging, while having epidermal concomitants, seems to involve primarily dermal and subcutaneous changes, and
25 is caused by (a) internal factors alone, as in intrinsic aging and (b) external factors, as in extrinsic aging. Intrinsic aging also is known as natural or chronologic aging, and extrinsic aging often is called photoaging. "Photodamage" implies skin damage caused by chronic sun exposure. These terms may be described as follows.

30 Intrinsic aging of skin, in sun-protected skin of the upper arm and abdomen, is an inherent degenerative process that occurs in all humans and is due to declining physiologic functions and capacities. This degenerative process may include qualitative and quantitative skin changes and also includes diminished or

defective synthesis of collagen and elastic fibers, and proteoglycans and glycosaminoglycans in the dermis. Signs of intrinsic aging include progressive thinning of skin, deepening of skin lines and fine wrinkles, lusterless skin surface, and loss of skin elasticity and recoilability. Although intrinsic aging of living
5 creatures is neither reversible nor preventable, modification and improvement of skin signs associated with such aging process can be achieved through topical management.

Extrinsic aging of skin is a distinctive process caused by external factors which include sunlight, radiation, air pollution, wind, cold, dampness, heat,
10 chemicals, smoke and cigarette smoking. Photoaging of skin may be defined as destructive cutaneous changes caused by chronic exposure to sunlight. Signs of photoaging on the face and back of hands include coarse and deepened wrinkles due to changes and degeneration of collagen and elastic fibers; marked loss of elasticity and recoilability; leathery skin surface and skin lesions with abnormal
15 pigmentation and increased numbers of age spots, pigmented spots, blotches and nodules. Histologically, the qualities and quantities of elastin and collagen tissues are changed. Normal elastin in tissues is replaced by abnormal elastin characterized as solar elastosis, and the normal collagen fibers are decreased.

Photodamage of skin, also called solar damage, may be defined as
20 cutaneous damage caused by chronic exposure to solar radiation and is associated with development of neoplastic lesions. Skin disorders caused by photodamage include pre-malignant lesions, basal cell carcinomas, squamous cell carcinomas and malignant melanomas.

There are numerous products available today that report to ameliorate some
25 of the signs of cutaneous aging. For example, there are a number of sunscreen patents related to treating aging skin. U.S. Patent No. 5,093,109 describes the use of titanium dioxide as a sunscreen which also contains antioxidants, emulsifiers, thickeners and colorants. U.S. Patent No. 5,204,105 describes the use of plant and yeast extracts, vitamins E & C to improve elasticity, the use of silicones to
30 improve firmness and the use of sunscreens to protect the skin from the sun. A number of cosmetic compositions are known that contain titanium dioxide either alone, or in combination with a silicone compound for various uses. See, U.S. Patent Nos. 4,801,445, 4,820,508, 5,032,390 and 5,250,289.

Other known compositions containing sunscreens together with free radical scavengers are described in, for example, U.S. Patent No. 5,093,109. These compositions, however, generally are oil in water emulsions containing ascorbyl palmitate, which rapidly falls apart in such an emulsion. U.S. Patent No. 5,560,917 describes a composition containing a sunscreen agent, a free radical scavenger and a cosmetically acceptable pigment.

There also are numerous descriptions in the literature of the use of alpha hydroxyacids and polyhydroxyacids for treating various skin disorders. U.S. Patent No. 3,879,537 describes the use of topical compositions containing an alpha hydroxyacid to alleviate the symptoms of ichthyosis. U.S. Patent No. 3,920,835 describes the use of topical compositions containing an alpha hydroxyacid to alleviate the symptoms of acne. U.S. Patent No. 3,984,566 describes the use of topical compositions containing an alpha hydroxyacid to improve the symptoms of dandruff.

Other document, for example, U.S. Patent Nos. 4,105,783, 4,197,316 and 4,380,549 describe the use of topical compositions containing an alpha hydroxyacid to alleviate or improve the symptoms of dry skin. In addition, U.S. Patent No. 4,234,599 describes the use of topical compositions containing an alpha hydroxyacid to alleviate the symptoms of actinic or nonactinic skin keratoses. Moreover, U.S. Patent No. 4,363,815 describes the use of topical compositions containing certain alpha hydroxyacids to improve skin conditions characterized by inflammation or disturbed keratinization.

Recently issued U.S. Patent Nos. 5,385,938 and 5,389,677 describe the use of compositions containing glycolic acid for topical treatment of skin wrinkles. Albert M. Kligman also described in U.S. Patent No. 4,877,805 that photoaging or sun damaged skin includes loss of collagen fibers, abnormal changes in elastic fibers and deterioration of small blood vessels in the dermis of the skin. The dermal components which make up the fibers of the dermis become smaller and sparser with increasing age, usually in sundamaged facial skin. There is a great loss of collagen fibers resulting in looseness and easy stretchability of the skin; elastic fibers become abnormal so that the skin does not promptly snap back after being stretched. Kligman disclosed the use of retin-A to treat photoaged or sundamaged skin.

These known compositions, are not always effective against the external effects of sunlight and the internal effects caused by intrinsic aging.

SUMMARY OF THE INVENTION

5 There exists a need for a single cosmetic composition that is effective not only against the external effects of sunlight but the internal effects caused by intrinsic aging. It is therefore an object of the present invention to provide such a composition. It is an additional object of the present invention to provide a composition that achieves the objectives described above, and at the same time
10 colors the skin attractively.

 In accordance with these and additional objects of the invention, there is provided a water in silicone emulsified cosmetic makeup composition including:

- 1) At least one hydroxy acid selected from polyhydroxy acid, alpha hydroxy acid, beta hydroxyacid, keto carboxylic acid and related
15 compounds;
- 2) Sunscreen agent(s);
- 3) Moisturizing agent(s);
- 4) Antioxidant(s);
- 5) Cosmetically acceptable pigment(s);
- 20 6) Silicone-containing compound(s); and
- 7) water.

 In accordance with additional objects of the invention, there are provided methods of reducing the appearance of lines and wrinkles, protecting the skin from harmful effects of ultraviolet light, moisturizing, soothing and firming the
25 skin and attractively coloring the skin comprising topically applying the above water in silicone emulsified cosmetic makeup composition in a cosmetically effective amount to achieve the aforementioned effects.

 In accordance with additional objects of the invention, there are provided methods of making a water in silicone emulsified cosmetic makeup composition
30 by:

- A) dissolving in water at least one hydroxy acid selected from polyhydroxy acid, alpha hydroxy acid, beta hydroxyacid, keto carboxylic acid and

related compounds in water and neutralizing to a pH of above 3.0 to provide phase A;

5 B) providing an aqueous mixture containing at least one moisturizing agent, heating the mixture, and adding this mixture to phase A to provide phase AB;

C) mixing at least one sunscreen agent, at least one antioxidant, at least one surfactant and at least one silicone-containing compound at room temperature, followed by heating and homogenizing the mixture to provide phase C;

10 D) providing a mixture containing at least one sunscreen agent and at least one cosmetically acceptable pigment, homogenizing the mixture to provide phase D;

E) mixing phases C) and D) together and homogenize until uniform to provide mixture CD;

F) slowly adding phase AB to phase CD to produce phase F;

15 G) adding to phase F a mixture containing parabens; and

H) adding additional cosmetically acceptable pigments to provide the appropriate color.

20 Additional objects and advantages of the invention will be set forth in part in the description that follows, and in part will be obvious from the description, or may be learned by practice of the invention. The objects and the advantages of this invention may be realized and obtained by means of the compositions and methods particularly pointed out in the appended claims.

DETAILED DESCRIPTION OF PREFERRED EMBODIMENT


25 Throughout this description, the expressions "alpha hydroxy acid," "AHA" and "2-hydroxycarboxylic acid" are used synonymously and denote any known alpha hydroxyacid useful in treating skin disorders, including those described in, *inter alia*, U.S. Patent No. 5,091,171, the disclosure of which is incorporated herein by reference in its entirety. Throughout this description, the expression
30 "polyhydroxy acid" denotes any of the known poly hydroxyacids reported as being useful in treating skin disorders, including poly hydroxyacids disclosed in U.S. Patent No. 5,091,171. Throughout this description, the expression "beta hydroxyacid" denotes any beta hydroxyacid useful in treating skin disorders.

Throughout this description, the expression "keto carboxylic acid" denotes any of the known keto carboxylic acids reported as being useful in treating skin disorders, including those disclosed in U.S. Patent No. 5,091,171. Throughout this description, the expression "related compound" refers to compounds that are structurally and functionally similar to alpha hydroxyacids and polyhydroxy acids such as those related compounds disclosed in U.S. Patent No. 5,091,171.

For convenience, the alpha hydroxyacids, polyhydroxy acids and related compounds which may be used in accordance with this invention may be classified into three groups, namely (1) 2-hydroxycarboxylic acids, (2) 2-ketocarboxylic acids and esters thereof, and (3) other related compounds. The related compounds may include hydroxycarboxylic acids with the hydroxyl group at any position other than position 2, for example position 3, position 4 or position 5, as well as cyclic hydroxycarboxylic acids (e.g., ascorbic acid and quinic acid), and also may include ketocarboxylic acids and esters thereof. Preferred related compounds include 3-hydroxycarboxylic acids, and 2-ketocarboxylic acids and esters thereof.

Group 1

The first group comprises organic carboxylic acids in which one hydroxy group is attached to the 2 position carbon atom of the acid. The generic structure of such 2-hydroxycarboxylic acids may be represented as follows:




Where R_a and R_b may be the same or different and are independently selected from H, F, Cl, Br, alkyl, aralkyl or aryl group of saturated or unsaturated, isomeric or non-isomeric, straight or branched chain or cyclic form, having 1 to 29 carbon atoms, and in addition R_a and R_b may carry OH, CHO, COOH and alkoxy group having 1 to 9 carbon atoms. In addition, the hydrogen atoms of R_a and/or R_b may be substituted by a nonfunctional element selected from Cl, Br, I, S, F, or a radical such as a low alkyl or alkoxy group, saturated or unsaturated, having 1 to 9 carbon atoms. The alpha hydroxy acids of Group 1 may be present as a free acid or lactone form, or in a salt form with an organic base or an inorganic alkali. The alpha hydroxy acids of Group 1 may exist as stereoisomers as D, L, and DL forms when R_a and R_b are not identical.


Typical alkyl, aralkyl and aryl groups for R_3 and R_5 include methyl, ethyl, propyl, isopropyl, butyl, pentyl, octyl, decyl, dodecyl, hexadecyl, benzyl, and phenyl, etc. Alpha hydroxy acids of Group 1 may be further divided into subgroups comprising (1) alkyl hydroxycarboxylic acids, (2) aralkyl and aryl hydroxycarboxylic acids, (3) polyhydroxy acids, and (4) hydroxy-polycarboxylic acids. The following are representative alpha hydroxy acids in each subgroup.

(1) Alkyl Hydroxycarboxylic Acids

1. 2-Hydroxyethanoic acid (Glycolic acid, hydroxyacetic acid)




2. 2-Hydroxypropanoic acid (Lactic acid)




3. 2-Methyl 2-hydroxypropanoic acid (Methylactic acid)




4. 2-Hydroxybutanoic acid



5. 2-Hydroxypentanoic acid




6. 2-Hydroxyhexanoic acid



7. 2-Hydroxyheptanoic acid



8. 2-Hydroxyoctanoic acid



9. 2-Hydroxynonanoic acid



10. 2-Hydroxydecanoic acid



11. 2-Hydroxyundecanoic acid



12. 2-Hydroxydodecanoic acid (Alpha hydroxylauric acid)



13. 2-Hydroxytetradecanoic acid (Alpha hydroxymyristic acid)

- (C₁₂H₂₅) (H) C (OH) COOH
14. 2-Hydroxyhexadecanoic acid (Alpha hydroxypalmitic acid)
(C₁₄H₂₉) (H) C (OH) COOH
15. 2-Hydroxyoctadecanoic acid (Alpha hydroxystearic acid)
(C₁₆H₃₃) (H) C (OH) COOH
16. 2-Hydroxyeicosanoic acid (Alpha hydroxyarachidonic acid)
(C₁₈H₃₇) (H) C (OH) COOH
17. 2-Hydroxytetraeicosanoic acid (Cerebronic acid)
(C₂₂H₄₅) (H) C (OH) COOH
18. 2-Hydroxytetraeicosenoic acid (Alpha hydroxynervonic acid)
(C₂₂H₄₃) (H) C (OH) COOH

(2) Aralkyl And Aryl 2-Hydroxycarboxylic Acids

1. 2-Phenyl 2-hydroxyethanoic acid (Mandelic acid)
(C₆H₅) (H) C (OH) COOH
2. 2,2-Diphenyl 2-hydroxyethanoic acid (Benzilic acid)
(C₆H₅) (C₆H₅) C (OH) COOH
3. 3-Phenyl 2-hydroxypropanoic acid (Phenyllactic acid)
(C₆H₅CH₂) (H) C (OH) COOH
4. 2-Phenyl 2-methyl 2-hydroxyethanoic acid (Atrolactic acid)
(C₆H₅) (CH₃) C (OH) COOH
5. 2-(4'-Hydroxyphenyl) 2-hydroxyethanoic acid
(4-Hydroxymandelic acid)
(HO-C₆H₄) (H) C (OH) COOH
6. 2-(4'-Chlorophenyl) 2-hydroxyethanoic acid
(4-Chloromandelic acid)
(Cl-C₆H₄) (H) C (OH) COOH
7. 2-(3'-Hydroxy-4'-methoxyphenyl) 2-hydroxyethanoic acid
(3-Hydroxy-4-methoxymandelic acid)
(HO-,CH₃O-C₆H₃) (H) C (OH) COOH

8. 2-(4'-Hydroxy-3'-methoxyphenyl) 2-hydroxyethanoic acid
(4-Hydroxy-3-methoxymandelic acid)
(HO-,CH₃O-C₆H₃) (H) C (OH) COOH
9. 3-(2'-Hydroxyphenyl) 2-hydroxypropanoic acid
[3-(2'-Hydroxyphenyl) lactic acid]
(HO-C₆H₄-CH₂) (H) C (OH) COOH
10. 3-(4'-Hydroxyphenyl) 2-hydroxypropanoic acid
[3-(4'-Hydroxyphenyl) lactic acid]
(HO-C₆H₄-CH₂) (H) C (OH) COOH
11. 2-(3',4'-Dihydroxyphenyl) 2-hydroxyethanoic acid
(3,4-Dihydroxymandelic acid)
(HO-,HO-C₆H₃) (H) C (OH) COOH
- (3) **Polyhydroxy Acids**
 1. 2,3-Dihydroxypropanoic acid (Glyceric acid)
(HOCH₂) (H) C (OH) COOH
 2. 2,3,4-Trihydroxybutanoic acid (Isomers; erythronic acid, threonic acid)
(HOCH₂ HOCH) (H) C (OH) COOH
 3. 2,3,4,5-Tetrahydroxypentanoic acid (Isomers; ribonic acid, arabinoic acid,
xylonic acid, lyxonic acid)
(HOCH₂ HOCH HOCH) (H) C (OH) COOH
 4. 2,3,4,5,6-Pentahydroxyhexanoic acid (Isomers; allonic acid, altronic acid,
gluconic acid, mannoic acid, gulonic acid, idonic acid, galactonic acid,
talonic acid)
(HOCH₂ HOCH HOCH HOCH) (H) C (OH) COOH
 5. 2,3,4,5,6,7-Hexahydroxyheptanoic acid (Isomers; glucoheptonic acid,
galactoheptonic acid etc.)
(HOCH₂ HOCH HOCH HOCH HOCH) (H) C (OH) COOH
- (4) **Hydroxy-polycarboxylic Acids**
 1. 2-Hydroxypropane-1,3-dioic acid (Tartronic acid)
(HOOC) (H) C (OH) COOH

2. 2-Hydroxybutane-1,4-dioic acid (Malic acid)
(HOOC CH₂) (H) C (OH) COOH
3. 2,3-Dihydroxybutane-1,4-dioic acid (Tartaric acid)
(HOOC HOCH) (H) C (OH) COOH
- 5 4. 2-Hydroxy-2-carboxypentane-1,5-dioic acid (Citric acid)
(HOOC CH₂)₂ C (OH) COOH
5. 2,3,4,5-Tetrahydroxyhexane-1,6-dioic acid (Isomers; saccharic acid, mucic acid etc.)
HOOC (CHOH)₄ COOH

10

The alpha hydroxy acids of Group 1 may be present in forms other than the acid, such as, for example, salts or lactones. Typical lactone forms which may be used in accordance with this invention include, for example, gluconolactone, galactonolactone, glucuronolactone, galacturonolactone, 15 gulonolactone, ribonolactone, saccharic acid lactone, pantoyllactone, glucoheptonolactone, mannonolactone, and galactoheptonolactone.

Group 2

The second group, which comprises compounds related to the alpha hydroxyacids of Group 1, includes organic carboxylic acids in which one keto group is attached to position 2 carbon atom of the acid. These compounds are referred to herein as keto carboxylic acids or 2-ketoacids. The generic structure of such 2-ketoacids, or alpha ketoacid may be represented as follows:



25

wherein R_c and R_d can be the same or different and are each selected from H, alkyl, aralkyl or aryl group of saturated or unsaturated, isomeric or non-isomeric, straight or branched chain or cyclic form, having 1 to 29 carbon atoms, and in addition R_c may carry F, Cl, Br, I, OH, CHO, COOH and alkoxy group having 1 to 9 carbon atoms. In addition, the hydrogen atoms of R_c and/or R_d may be substituted by a nonfunctional element selected from Cl, Br, I, S, F, or a radical 30 such as a low alkyl or alkoxy group, saturated or unsaturated, having 1 to 9 carbon atoms. The alpha ketoacids may be present as a free acid or an ester form, or in a salt form with an organic base or an inorganic alkali. The typical

alkyl, aralkyl and aryl groups for R_c and R_d include methyl, ethyl, propyl, 2-propyl, butyl, pentyl, hexyl, octyl, dodecyl, hexadecyl, benzyl and phenyl.

The ester form of the above-mentioned 2-ketoacids also is therapeutically effective for alleviating the signs and symptoms of cutaneous aging including intrinsic and extrinsic aging. For example, methyl 2-ketopropanoate and ethyl 2-ketopropanoate are therapeutically very effective. While the mechanism of action of these 2-ketoacids is not known, it is believed that the ester form of the 2-ketocarboxylic acid is chemically and/or biochemically very reactive, and a free 2-ketoacid may be released in the skin after penetration through the stratum corneum of the skin.

The representative 2-ketoacids and their esters of the second group are listed below:

1. 2-Ketoethanoic acid (Glyoxylic acid)
(H) CO COOH
2. Methyl 2-ketoethanoate
(H) CO COOCH₃
3. 2-Ketopropanoic acid (Pyruvic acid)
CH₃ CO COOH
4. Methyl 2-ketopropanoate (Methyl pyruvate)
CH₃ CO COOCH₃
5. Ethyl 2-ketopropanoate (Ethyl pyruvate)
CH₃ CO COOC₂H₅
6. Propyl 2-ketopropanoate (Propyl pyruvate)
CH₃ CO COOC₃H₇
7. 2-Phenyl-2-ketoethanoic acid (Benzoylformic acid)
C₆H₅ CO COOH
8. Methyl 2-phenyl-2-ketoethanoate (Methyl benzoylformate)
C₆H₅ CO COOCH₃
9. Ethyl 2-phenyl-2-ketoethanoate (Ethyl benzoylformate)
C₆H₅ CO COOC₂H₅
10. 3-Phenyl-2-ketopropanoic acid (Phenylpyruvic acid)
C₆H₅CH₂ CO COOH
11. Methyl 3-phenyl-2-ketopropanoate (Methyl phenylpyruvate)


- $C_6H_5CH_2 CO COOCH_3$
 12. Ethyl 3-phenyl-2-ketopropanoate (Ethyl phenylpyruvate)
 $C_6H_5CH_2 CO COOC_2H_5$
 13. 2-ketobutanoic acid
 5 $C_2H_5 CO COOH$
 14. 2-Ketopentanoic acid
 $C_3H_7 CO COOH$
 15. 2-Ketohexanoic acid
 $C_4H_9 CO COOH$
 10 16. 2-Ketoheptanoic acid
 $C_5H_{11} CO COOH$
 17. 2-Ketooctanoic acid
 $C_6H_{13} CO COOH$
 18. 2-Ketododecanoic acid
 15 $C_{10}H_{21} CO COOH$
 19. Methyl 2-ketooctanoate
 $C_6H_{14} CO COOCH_3$

Group 3

- 20 The third group, which also comprises related compounds, includes, *inter alia*, hydroxycarboxylic acids where the hydroxy is at a position other than position 2, and cyclic hydroxycarboxylic acids which are useful for topical application to improve signs of aging skin and the cutaneous appendages. The members of this group, which are more conveniently identified by name than by
 25 generic structures, include ascorbic acid, quinic acid, isocitric acid, tropic acid (2-phenyl 3-hydroxypropanoic acid), trethocanic acid, 3-chlorolactic acid, citramalic acid, agaricic acid, aleuritic acid, pantoic acid, lactobionic acid and hexulosonic acid.

30 Group 4

The fourth group includes beta hydroxyacids in which a hydroxy group is attached to the beta carbon atom of the acid. The generic structure of such beta hydroxyacids may be represented as follows:



where R_e , R_f are selected from the group consisting of hydrogen, alkyl, aralkyl and aryl, where said alkyl, aralkyl or aryl can be saturated or unsaturated, straight, branched or cyclic, and have from 1-25 carbon atoms. In addition, the hydrogen atoms of R_e and/or R_f may be substituted by a nonfunctional element selected from Cl, Br, I, S, F, or a radical such as a low alkyl or alkoxy group, saturated or unsaturated, having 1 to 9 carbon atoms. In the generic structure above, m preferably is an integer of from 1 to 9. The beta hydroxyacids may be present as a free acid or an ester form, or in a salt form with an organic base or an inorganic alkali. Preferably, R_e and R_f are selected from methyl, ethyl, propyl, 2-propyl, butyl, pentyl, hexyl, octyl, dodecyl, hexadecyl, benzyl and phenyl.

The alpha hydroxyacids, polyhydroxyacids, 2-ketoacids, beta hydroxyacids and related compounds are used in a cosmetically effective amount to reduce or alleviate the signs of fine lines and wrinkles. Preferably, these compounds are used in an additive amount within the range of from about 0.1 to about 25% by weight, based on the total weight of the composition. "Additive amount," as it is used herein denotes the total amount of component used in the composition, such as the total amount of alpha hydroxyacids, polyhydroxyacids, 2-ketoacids and related compounds used. More preferably, these compounds are used in an amount within the range of from about 2 to about 10% by weight and more preferably, from about 4 to about 8% by weight.

Preferably, the alpha hydroxyacids, polyhydroxyacids, 2-ketoacids and related compounds are neutralized with an appropriate base to a pH of more than about 3, more preferably more than about 3.5 and most preferably, to a pH of about 3.8. Any cosmetically acceptable base can be used to neutralize the alpha hydroxyacids, polyhydroxyacids, 2-ketoacids and related compounds. Preferably, the base is selected from sodium hydroxide or ammonium hydroxide or triethanolamine and most preferably, the base is ammonium hydroxide. Those skilled in the art are capable of determining an effective amount of alpha hydroxyacids, polyhydroxyacids, 2-ketoacids and related compound together with an acceptable quantity of base using the guidelines provided herein.

Throughout this description, the expression "sunscreen agent" denotes known sunscreen agents that are useful in absorbing, screening or preventing ultraviolet rays from penetrating the skin. Preferably, the sunscreen agent is titanium dioxide or zinc oxide, and more preferably the sunscreen agent is coated with a polymeric material or other cosmetically acceptable coating. Preferred sunscreen agents can be selected from microfine titanium dioxide coated with aluminum hydroxide and stearic acid (aluminum stearate), cosmetic grade titanium dioxide coated with C9-C15 polyfluoroalkyl phosphate, polymer coated zinc oxide, octyl dimethyl PABA, PABA and its esters, benzophenone-3, octyl salicylate, menthyl anthranilate, and phenylbenzimidazole sulfonic acid.

The sunscreen agent can be used in any effective amount that will prevent or inhibit ultraviolet rays from penetrating the skin. Specifically, the sunscreen agent is provided in an amount to provide a sunscreen protection factor of at least 8, more preferably, at least 15, and even more preferably, at least 30. Preferably, the sunscreen agent is used in an amount within the range of from about .1% to about 50% by weight, based on the total weight of the composition. More preferably, the sunscreen agent is used in an additive amount within the range of from about 0.1 to about 20% by weight, and most preferably, within the range of from about 6 to about 16% by weight.

Throughout this description, the expression "moisturizing agent(s)" denotes any agents that provide a moisturizing effect to the skin, e.g., any known humectant. Any moisturizing agent can be used in any moisturizing effective amount. Preferably, the moisturizing agent is selected from glycerin, butylene glycol or their derivatives such as 1,3-butylene glycol and glycerine, propylene glycol, sorbitol, sodium PCA, glucam E-10, glucam E-20, and POE (7-26) glycerine. Those skilled in the art recognize useful moisturizing agents that can be used in the present invention, using the guidelines provided herein. Preferably, the moisturizing agent is used in an amount within the range of from about 0.1 to about 10% by weight, based on the total weight of the composition, more preferably, from about 0.1 to about 5% by weight, and most preferably, from about 1 to about 5% by weight.

Any antioxidant(s) can be used in any effective amount to prepare the cosmetic composition of the invention. Preferably, the antioxidant is a natural

antioxidant such as a Vitamin E derivative, *e.g.*, Vitamin E linoleate, Vitamin E, Vitamin E POE succinate, Vitamin E acetate, ascorbic acid, ascorbyl palmitate and ascorbyl-PMG. The antioxidant useful in the invention can be used in an additive amount within the range of from about 0.1 to about 10% by weight, based on the total weight of the composition, preferably, from about 0.1 to about 5% by weight, and more preferably, from about 0.5 to about 1% by weight.

Throughout this description, the expression "cosmetically acceptable pigment" denotes any pigments that are useful in providing color to a cosmetic composition, and which do not adversely affect the skin. Any cosmetically acceptable pigment can be used in any effective amount to prepare the cosmetic composition of the invention. Preferably, the cosmetically acceptable pigment is an inorganic pigment such as iron oxide (yellow, red and black), manganese oxide, and F D & C Aluminum Lakes. Zinc oxide and titanium dioxide, although described herein as being useful sunscreen agents, can also be used within the context of the present invention as a pigment. The pigments, although useable in raw form as an iron oxide, preferably are coated with a polymeric coating, more preferably, the pigments are polyfluoroalkyl coated inorganic pigments. The pigments useful in the invention can be used in an additive amount within the range of from about 0.1 to about 25% by weight, based on the total weight of the composition, preferably, from about 1 to about 20% by weight, and more preferably, from about 2 to about 15% by weight.

The water in silicone emulsified cosmetic makeup composition of the present invention also includes silicone-containing compounds. Any known cosmetically silicone-containing compound can be used in a cosmetically effective amount. Preferably, the silicone-containing compound(s) is selected from cyclomethicone, dimethicone and derivatives thereof, such as cyclomethicone/quaternium-18 hectorite, cyclomethicone polyol, dimethicone polyol, and cetyl dimethicone copolyol and combinations thereof. Those skilled in the art will appreciate that the silicone-containing compound(s) also can function as an emulsifier, especially when the silicone-containing compound is selected from dimethicone polyol or cetyl dimethicone copolyol. Silica also is preferably used in the present invention. More preferably, the silicone-containing compounds are

selected from cyclomethicone, dimethicone, phenyl methicone, phenyl dimethicone and phenyl trimethicone.

The silicone-containing compounds preferably are used in an amount within the range of from about 1 to about 50% by weight, based on the total weight of the composition. More preferably, the silicone-containing compounds are used in an additive amount within the range of from about 5 to about 30% by weight, and most preferably, from about 10 to about 25% by weight.

The balance of the composition is comprised of customary additives selected from auxiliaries, filler, emulsifiers, surfactants, solvents, buffers, emollients and other cosmetically acceptable carriers and fillers, and water. Other conventional additives can be used in the present invention, including antioxidants, dispersants, humectants, thickeners and preservatives. Water typically is deionized water, and the amount of water typically is anywhere from about 10 to about 80 by weight, based on the total weight of the composition, preferably from about 15 to about 75% by weight, and most preferably from about 20 to about 40% by weight. The amount of conventional additives typically ranges from about 0 to about 20% by weight, based on the total weight of the composition, preferably from about 1 to about 15% by weight, and most preferably from about 4 to about 10% by weight.

As an emulsion, the emulsified composition of the present invention includes an oil phase and an aqueous phase. The oil phase preferably is present in an amount within the range of from about 1 to about 75% by weight, based on the total weight of the composition, more preferably, from about 5 to about 60% by weight, and most preferably from about 40 to about 60% by weight.

Emulsifiers can be used to assist in preparing the emulsified cosmetic makeup composition of the invention. Any cosmetically acceptable emulsifier can be used in an amount that provides the desired emulsifying effect. Preferably, the emulsifier is selected from known soaps and surfactants, more preferably, the emulsifier is selected from stearic acid, sorbitan sesquioleate, polyethylene glycol (PEG)-30 dipolyhydroxystearate, lecithin, magnesium stearate, and derivatives and mixtures thereof. The emulsifiers preferably are used in an additive amount within the range of from about 0.5 to about 30% by weight, based on the total

weight of the composition, more preferably, from about 1 to about 12% by weight and most preferably from about 4 to about 8% by weight.

Those skilled in the art are capable of selecting particular ingredients for use in the present composition, and they are capable of determining an appropriate amount of ingredient to provide a water in silicone emulsified cosmetic makeup composition.

A particularly preferred composition for reducing the appearance of fine lines and wrinkles, moisturizing and protecting the skin from the harmful effects of sunlight includes a water in silicone emulsion containing about 0.1 to 25 weight % ultrafine titanium dioxide, from about 0.1 to 25 weight % of microfine zinc oxide, from about 0.1 to 25 weight % of a mixture of gluconolactone (as a polyhydroxy acid) and glycolic acid (as an alpha hydroxyacid) neutralized to a pH of 3.80 with any base, preferably ammonium hydroxide, from about 0.1 to 5 weight % of vitamin E derivative, preferably vitamin E linoleate, from about 0.1 to 5 weight % of a moisturizing agent, preferably glycerin and from about 0.1 to 25% of a blend of cosmetically acceptable pigments, preferably polyfluoroalkyl coated inorganic pigments.

Even more preferred embodiments of the invention include the use of microfine titanium dioxide coated with aluminum hydroxide and stearic acid (aluminum stearate), cosmetic grade titanium dioxide coated with C9-C15 polyfluoroalkyl phosphate, and polymer coated zinc oxide as sunscreen agents, Gluconolactone and/or glycolic acid as the hydroxy acid(s), Vitamin E linoleate as a natural antioxidant, glycerin and/or butylene glycol as moisturizing agent(s) (*i.e.*, humectant(s)), and cosmetically acceptable iron oxide pigments.

To prepare an emulsified cosmetic composition in accordance with the present invention, at least one hydroxy acid or related compound first is dissolved in a solution prepared from ethanol, water, propylene glycol, butylene glycol, acetone or other pharmaceutically acceptable vehicle, and then neutralized with an appropriate base. The emulsified cosmetic composition can be prepared as follows:

A) dissolving in water at least one hydroxy acid selected from polyhydroxy acid, alpha hydroxy acid, beta hydroxy acid, keto carboxylic acid

and related compounds in water and neutralizing with an appropriate base to a pH of above 3.0 to provide phase A;

5 B) providing an aqueous mixture containing at least one moisturizing agent, heating the mixture, and adding this mixture to phase A to provide phase AB;

C) mixing at least one sunscreen agent, at least one antioxidant, at least one emulsifier and at least one silicone-containing compound at room temperature, followed by heating and homogenizing the mixture to provide phase C;

10 D) providing a mixture containing at least one sunscreen agent and at least one cosmetically acceptable pigment, homogenizing the mixture to provide phase D;

E) mixing phases C) and D) together and homogenize until uniform to provide mixture CD;

F) slowly adding phase AB to phase CD to produce phase F;

15 G) adding to phase F a mixture containing parabens; and

H) adding additional cosmetically acceptable pigments to provide the appropriate color.

20 In step A), the hydroxyacid preferably is gluconolactone and/or glycolic acid. In step B), the moisturizing agent(s) preferably are 1,3-butylene glycol and glycerine and the mixture is heated to a temperature within the range of from about 35-50°C, preferably from about 40-45°C. The mixture of step B) preferably is stirred for about 10 minutes prior to addition to phase A). It is preferred to mix the ingredients of phase C) together at room temperature, whereby the pigment preferably is titanium dioxide, the emulsifier preferably is sorbitan sesquileate and lecithin, the antioxidant preferably is Vitamin E linoleate
25 and the silicone-containing compound preferably is selected from cyclomethicone and dimethicone. The components of phase C) preferably are mixed together at room temperature, followed by heating and homogenizing for about 30 minutes or until the mixture is substantially homogeneous.

30 In preparing the emulsified cosmetic composition of the invention, it is preferred to use only a portion of the cosmetically acceptable pigment to prepare phase D), more preferably, use only 90% of the pigment, most preferably 80%. The remaining pigment may be added later in step H) to adjust the color of the

cosmetic composition. The mixture of phase D) preferably is homogenized for about 30 minutes until uniform. Phase D) then is added to phase C) and the mixture of CD) is homogenized for about 30 minutes until uniform. In step F), phase AB) is slowly added to phase CD) with slow speed mixing. A mixture containing parabens then is added to the mixture of AB) and CD) with high mixing and mixed for at least 5 minutes, and the color of the final emulsion is adjusted using the remaining pigment from step D).

The present invention also encompasses a method of reducing the appearance of fine lines and wrinkles, protecting the skin from ultraviolet radiation exposure, moisturizing and soothing the skin comprising topically applying a water in silicone emulsified cosmetic makeup composition of the invention to an area of the skin in need thereof. Preferably, the composition is applied to areas of the skin that are typically exposed to ultraviolet radiation, such as the arms, hands, face, feet, legs, chest, buttocks and abdomen, although the composition of the invention can be applied to any area of skin. The composition of the present invention preferably is applied on a daily basis, and more preferably, twice daily in order to achieve the appropriate effects. Preferably, the composition of the present invention is applied twice daily for a period of at least 2 months, preferably at least 5 months, and most preferably more than 8 months to achieve a reduction in the appearance of fine lines and wrinkles. The rate and duration of application can be varied, however, depending on the concentration of effective ingredients. Skilled artisans are capable of designing a suitable application regimen to provide the requisite fine line and wrinkle reducing effect as well as ultraviolet exposure reducing effect using the guidelines provided herein.

The following are illustrative examples of formulations and compositions according to this invention. Although the examples utilize only selected compounds and formulations, it should be understood that the following examples are illustrative and not limited. Therefore, any of the known alpha hydroxyacids, polyhydroxy acids, beta hydroxyacids, keto carboxylic acids, related compounds, moisturizing agents, antioxidants, silicone-containing compounds, emulsifiers, pigments and sunscreen agents may be substituted according to the teachings of this invention in the following examples.

EXAMPLES

A preferred makeup composition and its method of manufacture is described below.

	PHASE	INGREDIENTS	%
5	A	DI WATER	20.900
		GLUCONOLACTONE	4.000
		AMMONIUM HYDROXIDE	0.600
		SODIUM CHLORIDE	1.000
	B	1,3-BUTYLENE GLYCOL	3.000
		PEG-30 DIPOLYHYDROXYSTEARATE	1.000
		N-ETHYL N-SOYAMORPHOLINIUM	1.000
		ETHOSULFATE/WATER	
		GLYCERINE	2.000
	C	TITANIUM DIOXIDE/ISOSTEARYL	10.000
		NEOPENTANOATE/	
		STEARIC ACID/ALUMINUM	
		HYDROXIDE	
		POLYMETHYL METHACRYLATE	2.000
		CYCLOMETHICONE/DIMETHICONE	5.000
		COPOLYOL	
		NYLON-12	2.500
		ISODODECANE	3.500
		ISOHEXADECANE	3.000
		VITAMIN E LINOLEATE	0.500
		SORBITAN SESQUIOLEATE	1.000
		CETYL DIMETHICONE COPOLYOL	1.500
		CYCLOMETHICONE/QUATERNIUM-18	5.000
		HECTORITE/	
		PROPYLENE CARBONATE	
		LECITHIN	0.500
		MAGNESIUM STEARATE	0.500
		POLYETHYLENE	2.000
		SILICA	1.500
		SILK POWDER	1.000
	D	TITANIUM DIOXIDE	3.000
		CYCLOMETHICONE	15.000
		TALC	0.500
		IRON OXIDE YELLOW	5.330
		IRON OXIDE RED	1.420
		IRON OXIDE BLACK	0.750
	E	PHENOXYETHANOL/ISOPROPYL	1.000
		PARABEN/	
		ISOBUTYL PARABEN/BUTYL PARABEN	
			100.00

The above cream makeup composition was prepared as follows, whereby the amounts of ingredients are indicated above in weight percent.

1) Gluconolactone was dissolved in water, stirred for about 3 hours, and the pH was adjusted to 3.80 with addition of ammonium hydroxide. Sodium chloride then was added to the solution to prepare phase A.

2) The ingredients for phase B then were mixed together and heated to 40-45°C. The heated mixture was stirred for about 10 minutes and added to phase A above to provide phase AB.

3) The ingredients of phase C were mixed together in a separate container at room temperature, followed by heating and homogenizing for 30 minutes until uniform to provide phase C.

4) In a container separate from that used to prepare phase C), and using only 80% of the iron oxide pigment, the ingredients of phase D (less the 20% of iron oxide pigment) were mixed together and homogenized for 30 minutes until uniform to provide phase D.

5) Phase D then was added to phase C to prepare phase CD), and phase CD) was homogenized for 30 minutes until uniform.

6) The ingredients of phase AB then were slowly added to phase CD with slow speed mixing.

7) The ingredients of phase E then were added to the emulsion of step 6), and the emulsion was homomixed at high speed for 5 minutes. The color of the resulting emulsified cosmetic composition was adjusted and matched using the remaining 20% of iron oxide pigments from step 4) to prepare a water in silicone emulsified cosmetic makeup composition.

The water in silicone emulsified cosmetic makeup composition then can be applied to various places on the body, preferably on the face, arms and hands, and used as both an effective sunscreen and as a composition effective in removing or alleviating the signs of aging such as fine lines and wrinkles.

It will be apparent to those skilled in the art that various modifications and variations can be made to the compositions of matter and methods of this invention. Thus, it is intended that the present invention covers such modifications and variations.

What is Claimed Is:

1. A water in silicone emulsified cosmetic makeup composition comprising:

- 1) At least one hydroxy acid selected from polyhydroxy acid, alpha hydroxy acid, beta hydroxyacid, keto carboxylic acid and related compounds;
- 2) Sunscreen agent(s);
- 3) Moisturizing agent(s);
- 4) Antioxidant(s);
- 5) Cosmetically acceptable pigment(s);
- 6) Silicone-containing compound(s); and
- 7) water.

2. The makeup composition as claimed in claim 1, wherein the at least one hydroxy acid is gluconolactone.

3. The makeup composition as claimed in claim 1, wherein the alpha hydroxy acid is glycolic acid.

4. The makeup composition as claimed in claim 1, wherein the sunscreen agent is comprised of ultrafine particles of titanium dioxide coated with aluminum hydroxide and stearic acid.

5. The makeup composition as claimed in claim 1, wherein the sunscreen agent is comprised of ultrafine particles of zinc oxide coated with polymeric materials.

6. The makeup composition as claimed in claim 1, wherein the moisturizing agent(s) is selected from glycerin or butylene glycol.

7. The makeup composition as claimed in claim 1, wherein the antioxidant is Vitamin E linoleate.

8. The makeup composition as claimed in claim 1, wherein the cosmetically acceptable pigment is at least one C9-C15 polyfluoroalkyl treated iron oxide.

9. The makeup composition as claimed in claim 4, wherein the sunscreen agent is present in an amount sufficient to provide a sun protection factor of at least 15.

10. The makeup composition as claimed in claim 1, wherein the emulsifier is selected from one or more emulsifiers selected from the group consisting of dimethicone copolyol, sorbitan sesquioleate, PEG-30 dipolyhydroxystearate and cetyl dimethicone copolyol.

11. A method of reducing the appearance of fine lines and wrinkles, protecting the skin from harmful effects of ultraviolet light, moisturizing and soothing the skin comprising topically applying a water in silicone emulsified cosmetic makeup composition as claimed in claim 1, in a cosmetically effective amount and for an effective period of time.

12. A method of making a water in silicone emulsified cosmetic makeup composition as claimed in claim 1, comprising:

A) dissolving in water at least one hydroxy acid selected from polyhydroxy acid, alpha hydroxy acid, beta hydroxy acid, keto carboxylic acid and related compounds in water and neutralizing the solution to a pH of above 3.0 to provide phase A;

B) providing an aqueous mixture containing at least one moisturizing agent, heating the mixture, and adding this mixture to phase A to provide phase AB;

C) mixing in a separate container from phase AB, at least one sunscreen agent, at least one antioxidant, at least one surfactant and at least one silicone-containing compound at room temperature, followed by heating and homogenizing the mixture to provide phase C;

D) mixing in a container separate from phase AB and phase C, at least one sunscreen agent and at least one cosmetically acceptable pigment, and homogenizing the mixture to provide phase D;

- E) mixing phases C) and D) together and homogenizing the mixture until uniform to provide phase CD;
- F) slowly adding phase AB to phase CD to produce phase F;
- G) adding to phase F a mixture containing parabens; and
- H) adding additional cosmetically acceptable pigments to provide the appropriate color.

13. The method as claimed in claim 12, wherein the at least one hydroxy acid is gluconolactone.

14. The method as claimed in claim 12, wherein the alpha hydroxy acid is glycolic acid.

15. The method as claimed in claim 12, wherein the sunscreen agent is comprised of ultrafine particles of titanium dioxide coated with aluminum hydroxide and stearic acid.

16. The method as claimed in claim 12, wherein the sunscreen agent is comprised of ultrafine particles of zinc oxide coated with polymeric materials.

17. The method as claimed in claim 12, wherein the moisturizing agent(s) is selected from glycerin or butylene glycol.

18. The method as claimed in claim 12, wherein the antioxidant is Vitamin E linoleate.

19. The method as claimed in claim 12, wherein the cosmetically acceptable pigment is at least one C9-C15 polyfluoroalkyl treated iron oxide.

20. The method as claimed in claim 12, wherein the emulsifier is selected from one or more emulsifiers selected from the group consisting of dimethicone copolyol, sorbitan sesquioleate, PEG-30 dipolyhydroxystearate and cetyl dimethicone copolyol.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 98/26580

A. CLASSIFICATION OF SUBJECT MATTER IPC 6 A61K7/48 A61K7/42 A61K7/021		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 6 A61K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 95 04517 A (THE PROCTER & GAMBLE COMPANY ET AL.) 16 February 1995 see page 20 - page 23 see claims 1,6-8,16,17,19,27,33	1,3,6,10
Y	---	4,5
Y	EP 0 796 612 A (L'OREAL) 24 September 1997 see page 3, line 35 - line 47 see page 4, line 4 - line 6 see page 4, line 53 - page 5, line 4 see page 5, line 15 --- -/--	4,5
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "&" document member of the same patent family		
Date of the actual completion of the international search 31 March 1999		Date of mailing of the international search report 08/04/1999
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer Alvarez Alvarez, C

INTERNATIONAL SEARCH REPORT

I. International Application No
PCT/US 98/26580

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 691 126 A (BEIERSDORF AG) 10 January 1996 see page 4, line 10 - page 5, line 58 see page 6, line 28 - page 7, line 7 see page 7, line 36 see page 7, line 45 - line 50 see page 7, line 57 - line 58 see page 8, line 1 - line 3 see example 1	1,2,9-11
A	---	12-20
X	EP 0 456 459 A (UNILEVER PLC ET AL.) 13 November 1991 see page 4, line 19 - line 24 see page 9, line 21 - line 52 see page 11 - page 14	1,4
A	---	
A	EP 0 808 625 A (KAO CORPORATION) 26 November 1997 see page 27, line 1 - line 30 see page 19, line 1 - line 30 -----	1

INTERNATIONAL SEARCH REPORT

Information on patent family members

In tional Application No

PCT/US 98/26580

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9504517 A	16-02-1995	AU 694740 B AU 7517594 A CA 2168842 A CN 1130861 A CZ 9600346 A EP 0717612 A JP 9501431 T US 5871754 A	30-07-1998 28-02-1995 16-02-1995 11-09-1996 12-06-1996 26-06-1996 10-02-1997 16-02-1999
EP 796612 A	24-09-1997	FR 2746301 A FR 2746302 A JP 10007520 A	26-09-1997 26-09-1997 13-01-1998
EP 691126 A	10-01-1996	DE 4423450 A JP 8026972 A	11-01-1996 30-01-1996
EP 456459 A	13-11-1991	AT 103170 T AU 636483 B AU 7640691 A CA 2041917 A DE 69101466 D DE 69101466 T DK 456459 T ES 2062683 T GB 2243780 A,B IN 172888 A JP 4226906 A JP 7045375 B US 5196187 A	15-04-1994 29-04-1993 14-11-1991 11-11-1991 28-04-1994 14-07-1994 25-07-1994 16-12-1994 13-11-1991 25-12-1993 17-08-1992 17-05-1995 23-03-1993
EP 808625 A	26-11-1997	NONE	